

Durham Research Online

Deposited in DRO:

08 March 2016

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Brown, T.J. and Todd, A. and O'Malley, C. and Moore, H.J. and Husband, A.K. and Bamba, C. and Kasim, A. and Sniehotta, F.F. and Steed, L. and Smith, S. and Nield, L. and Summerbell, C.D. (2016) 'Community pharmacy-delivered interventions for public health priorities : a systematic review of interventions for alcohol reduction, smoking cessation and weight management, including meta-analysis for smoking cessation.', *BMJ open.*, 6 (2). e009828.

Further information on publisher's website:

<http://dx.doi.org/10.1136/bmjopen-2015-009828>

Publisher's copyright statement:

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

BMJ Open Community pharmacy-delivered interventions for public health priorities: a systematic review of interventions for alcohol reduction, smoking cessation and weight management, including meta-analysis for smoking cessation

Tamara J Brown,^{1,2,3} Adam Todd,^{1,2,3,4} Claire O'Malley,^{1,2,3} Helen J Moore,^{1,2,3} Andrew K Husband,^{1,2,3} Clare Bamba,^{2,3,4} Adetayo Kasim,³ Falko F Sniehotta,^{2,5} Liz Steed,⁶ Sarah Smith,^{1,2,3} Lucie Nield,⁷ Carolyn D Summerbell^{1,2,3}

To cite: Brown TJ, Todd A, O'Malley C, *et al.* Community pharmacy-delivered interventions for public health priorities: a systematic review of interventions for alcohol reduction, smoking cessation and weight management, including meta-analysis for smoking cessation. *BMJ Open* 2016;**6**:e009828. doi:10.1136/bmjopen-2015-009828

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2015-009828>).

Received 26 August 2015
Revised 26 November 2015
Accepted 11 January 2016



CrossMark

For numbered affiliations see end of article.

Correspondence to
Professor Carolyn Summerbell;
carolyn.summerbell@durham.ac.uk

ABSTRACT

Objectives: To systematically review the effectiveness of community pharmacy-delivered interventions for alcohol reduction, smoking cessation and weight management.

Design: Systematic review and meta-analyses. 10 electronic databases were searched from inception to May 2014.

Eligibility criteria for selecting studies: *Study design:* randomised and non-randomised controlled trials; controlled before/after studies, interrupted times series. *Intervention:* any relevant intervention set in a community pharmacy, delivered by the pharmacy team. No restrictions on duration, country, age, or language.

Results: 19 studies were included: 2 alcohol reduction, 12 smoking cessation and 5 weight management. Study quality rating: 6 'strong', 4 'moderate' and 9 'weak'. 8 studies were conducted in the UK, 4 in the USA, 2 in Australia, 1 each in 5 other countries. Evidence from 2 alcohol-reduction interventions was limited. Behavioural support and/or nicotine replacement therapy are effective and cost-effective for smoking cessation: pooled OR was 2.56 (95% CI 1.45 to 4.53) for active intervention vs usual care. Pharmacy-based interventions produced similar weight loss compared with active interventions in other primary care settings; however, weight loss was not sustained longer term in a range of primary care and commercial settings compared with control. Pharmacy-based weight management interventions have similar provider costs to those delivered in other primary care settings, which are greater than those delivered by commercial organisations. Very few studies explored if and how sociodemographic or socioeconomic variables moderated intervention effects. Insufficient information was available to examine relationships between effectiveness and behaviour change strategies,

Strengths and limitations of this study

- To the best of our knowledge, this is the first systematic review that combines evidence from community pharmacy-delivered alcohol, smoking and weight management interventions, and directly compares these findings with other primary care and community healthcare settings.
- This review provides healthcare commissioners with useful evidence on reach, effectiveness and costs when considering using community pharmacies to help deliver smoking cessation and weight management services.
- There was insufficient evidence to assess the effectiveness of community pharmacy-based interventions on health equity.
- The descriptions available did not allow for the coding of specific aspects of theory and behavioural content of the interventions.
- Insufficient information was available to examine the relationship between intervention effectiveness and behaviour change strategies and/or models used, implementation factors, or the organisation and delivery of interventions.

implementation factors, or organisation and delivery of interventions.

Conclusions: Community pharmacy-delivered interventions are effective for smoking cessation, and demonstrate that the pharmacy is a feasible option for weight management interventions. Given the potential reach, effectiveness and associated costs of these interventions, commissioners should consider using community pharmacies to help deliver public health services.

INTRODUCTION

A number of agencies and countries, including WHO, have set a clear strategy for the future of public health. This agenda is focused on improving the healthy life expectancy of the population and, where possible, reducing or removing threats to this aim.¹ One strand within this agenda is to create accessible, multidisciplinary networks of public health professionals who work within communities and provide services to address key public health issues, health inequalities, and ultimately improve health and well-being. Worldwide, community pharmacies may be an important component of this agenda; WHO acknowledges that community pharmacies and their staff are easily accessible and, as such, could play a key role in delivering public health initiatives, especially in priority areas.² For example, in England, community pharmacies are more accessible than general practice (GP) services.³ A recent study has also demonstrated that, in England, 89% of the population can walk to a community pharmacy within 20 min. Significantly, in areas of highest deprivation, this value increases to almost 100%—the so-called positive pharmacy care law.⁴ Community pharmacies could, therefore, be a way of engaging with hard-to-reach populations.

In view of this, many community pharmacies in some countries, now offer smoking cessation services, and a few offer alcohol and weight reduction services.⁵ These services are delivered by pharmacists, pharmacy technicians and/or medicine counter assistants, with a view to modifying health-related behaviours. The specific types of services are wide ranging and include two main approaches: pharmaceutical-related (eg, supplying nicotine replacement therapy (NRT), monitoring of biochemical markers) and non-pharmaceutical-related (eg, providing advice on behaviour change strategies), or a combination of both approaches. Funding arrangements for these services vary by country; in the UK, at present, many of these services are commissioned by the local authority according to local need, and delivered according to an agreed framework. Currently, six Local Pharmaceutical Committees (LPCs) have weight management services, 14 LPCs have alcohol reduction services, and there are 81 stop smoking services.⁶

In 2008, the Department of Health for England⁷ stated it was important to develop 'a sound evidence base that demonstrates how pharmacy delivers effective, high quality and value for money services'. Reviews published since 2008 have attempted to summarise this evidence base, but the lack of relevant randomised controlled trials (RCTs) have limited their findings.^{8–10} However, a scoping search performed in 2013 revealed a number of relevant RCTs that had, or were about to report their findings; a number of relevant controlled trials were also identified, that could usefully inform the evidence base where there was a lack of evidence from RCTs. The primary objective of this review, therefore, was to systematically review the effectiveness of community pharmacy-delivered interventions for alcohol

reduction, smoking cessation and weight management. The secondary objectives were to explore if and how age, ethnicity, gender and socioeconomic status (SES), moderate intervention effects; and to describe how the interventions have been implemented, organised and delivered.

METHODS

The review was funded by the National Institute for Health Research Public Health Research Programme (project number 12/153/52). The review was carried out using the principles outlined in the Cochrane Handbook for Systematic Reviews of Interventions.¹¹ The protocol is published in *BMC Systematic Reviews*,¹² and is registered with the International Prospective Register of Systematic Reviews (PROSPERO CRD42013005943). A review advisory group comprising patients, pharmacists and researchers, helped to guide the research. The review is reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.^{13 14}

Interventions

The review included any type of community pharmacy-delivered intervention aimed at alcohol reduction, smoking cessation, or weight management; of any duration, based in any country and of any age. The setting of interest was the community pharmacy, which was defined as a pharmacy set in the community, which is accessible to all and not based in a hospital, clinic or online. Where a pharmacy is referred to throughout this paper, we refer to a community pharmacy. There was no restriction on the type of comparator, which could be a non-active control, usual care, or another type of active intervention, set in or out of the community pharmacy. Participants could be recruited from outside of the community pharmacy as long as one of the intervention groups was delivered from the community pharmacy. The intervention had to be delivered by the community pharmacist, pharmacy technician or medicines counter assistant; however, the intervention could also include other deliverers as part of a multidisciplinary team.

Study design

A broad range of controlled study designs were included, using the Cochrane Effective Practice and Organisation of Care (EPoC) study design criteria.¹⁵ These included RCTs; non-RCTs (nRCT); controlled before/after studies (CBA); interrupted time series (ITS), and repeated measures studies. We included both fully powered and pilot studies; studies were graded lower on quality if they were insufficiently powered.

Search strategy

Ten electronic databases were searched: Applied Social Sciences Index and Abstracts, Cumulative Index to Nursing and Allied Health Literature, EMBASE,

International Bibliography of the Social Sciences, MEDLINE, NHS Economic Evaluation Database, PsycINFO, Social Science Citation Index, Scopus and the Sociological Abstracts; from inception to May 2014 (see online supplementary file 1). Supplementary searches to identify published, unpublished and ongoing studies included bibliographies, contacting experts, grey literature (OpenGrey, Social Care Online, Prevention Information & Evidence elibrary and Nexus UK), study registers (International Standard Registered Clinical/soCial sTudy Number registry and the National Research Register) and website (Google).

Outcomes

Interventions for alcohol reduction and smoking cessation had to report a relevant behavioural outcome, and interventions for weight had to report an anthropometric outcome. These outcomes were considered the primary outcomes, and could be measured or self-reported. Where studies reported if and how sociodemographic (age, ethnicity, gender) and/or SES (education, income, occupation, social class, deprivation or poverty) moderated intervention effects on the primary outcomes, this is reported in the review.

The review also describes how the interventions have been organised, implemented and delivered using the methodological tool for the assessment of the implementation of complex public health interventions in systematic reviews, developed by Egan *et al*¹⁶ for the workplace, and adapted by Bambra *et al*¹⁷ for obesity interventions. The Behaviour Change Wheel¹⁸ and the Nuffield Intervention Ladder¹⁹ were used to broadly describe the behavioural strategies, intervention functions and policy categories of the interventions.

Data extraction and quality appraisal

Three reviewers (CO, HM, SS) screened the titles and abstracts and two reviewers (CO, TB) screened the full-text articles. Data extraction and quality assessment were conducted independently by TB and one other reviewer (from among AT, CO, CS, HM, LN, LS, SS). Study quality was appraised using the Effective Public Health Practice Project Quality Assessment Tool for Quantitative Studies,²⁰ which is recommended by the Cochrane Public Health Review Group.²¹ Studies were assessed for quality using six criteria: selection bias, study design, confounders, blinding, data collection methods and withdrawals/dropouts. Each study was given an overall (global) rating based on the ratings for the six criteria: 'strong' (no 'weak' ratings), 'moderate' (one 'weak' rating) and 'weak' (two or more 'weak' ratings). Any discrepancies in the data extraction or quality assessment were resolved through discussion, or referred to a third reviewer (CS) for final assessment. Extraction of contextual data was conducted by one reviewer (CS) and checked by another (TB). Assessment of behaviour change strategies used was conducted by one reviewer (CS) and checked by two others (FS and LS).

Analysis and synthesis

Narrative synthesis was conducted for all the included interventions. Owing to the heterogeneity of the studies, it was only possible to conduct meta-analyses for the smoking cessation studies. The smoking data was analysed (AK) using binomial-normal random effect model (R package meta). In order to explain the observed heterogeneity between studies, four different meta-regression models were fitted, accounting for whether the comparator was an active control or usual care, duration of the intervention and the global quality assessment ratings. Q-statistics and the percentage of heterogeneity between studies were reported for each meta-regression model. The most optimal meta-regression model was chosen using a minimum Akaike Information criterion. Owing to the limited available data and lack of informative priors, subgroup analysis by demographic or SES was not considered. A funnel plot for the smoking cessation RCTs was carried out to indicate the possible presence of publication bias and other biases.

RESULTS

The electronic search identified over 19 000 records, of which 72 full-text articles were screened for eligibility; 19 studies (from 23 articles) were included, and 49 were excluded. Five excluded studies^{22–26} (from six articles) were pharmacotherapy plus lifestyle advice interventions in participants with comorbidities. These studies were excluded because the primary focus was not alcohol, smoking or weight management; these interventions focused on self-management of a chronic condition. The process of inclusion and exclusion of studies are shown in figure 1.

Study characteristics

Tables 1–3 provide the main study characteristics for all 19 interventions (see online supplementary file 2) for detailed study characteristics, including sociodemographic and SES). There were 2 alcohol reduction interventions,^{27 28} 12 smoking cessation interventions^{29–40} and 5 weight management interventions.^{41–45} There were 15 RCTs, 2 nRCTs^{25 33 44} and 2 CBAs.^{29 29 42} There were 17 published journal articles and two reports.^{28 42} Eight studies were conducted in the UK,^{27–29 33 36 38 42 43} four in the USA,^{30 35 41 44} two in Australia^{31 39} and one each in Canada,³² Denmark,⁴⁰ Japan,³⁷ The Netherlands³⁴ and Thailand.⁴⁵ All studies were of adults. Fourteen studies reported on funding; types of funding sources included academic research bodies, health-related institutions, commercial organisations and pharmaceutical companies.

Three studies^{29 42 43} recruited participants from areas of high deprivation, and compared a pharmacy-based setting with other settings. Twelve studies recruited participants within the community pharmacy; other recruitment settings included hospital/primary care units, via telephone and a community health centre. Types of




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram.

pharmacies included single outlets, small chains and large chains; set in rural, urban and a combination of both geographical settings. The number of pharmacies included within each study ranged from one to over 200. Participant sample size ranged from 28 to around 7000, comprising approximately 13 500 service users in total. Mean age ranged from 24 to 60 years; there was a majority of females across all studies, particularly in the weight management studies. Duration of follow-up ranged from 5 to 56 weeks.

In terms of data analysis, only four studies assessed whether sociodemographic variables moderated the effect of interventions; four studies^{28 38 42 43} assessed any differential effects of gender, and one of these also assessed age.⁴² No study assessed any differential effects of SES. Few studies used regression analysis to assess the influence of sociodemographic or socioeconomic variables on change from baseline, as potential predictors of outcomes within intervention groups, or to explain retention.

Quality assessment

The studies were assessed for quality using six criteria and assigned a global rating; six studies were rated 'strong', four studies 'moderate' and nine studies 'weak' (see online supplementary file 3). Participants were not obtained from a randomly selected sample in any of the studies. Five studies reported a low attrition rate, with follow-up of at least 80% of participants. Only five studies were sufficiently powered. Six studies conducted intention-to-treat analyses. Fifteen studies imputed data from baseline or last follow-up, or made assumptions

about dropouts (eg, assumed that dropouts had not stopped smoking/not lost weight). Six studies used hierarchical modelling techniques to adjust for potential pharmacy or pharmacist-level effects on individual participant outcomes. None of the studies reported details about whether the intervention was delivered as intended, for example, by observation of sessions, quality control audits, or staff and researcher records.

Implementation of the interventions

Very few studies reported any degree of consultation or collaboration, with stakeholders as part of the planning process, or during delivery of the intervention (see online supplementary file 4). Both the brief alcohol reduction interventions consulted with pharmacists during the planning stages.^{28 27} The smoking cessation study by Hoving *et al*³⁴ collaborated with a national charity on smoking and health, and together they developed the intervention. The smoking cessation study by Costello *et al*³² was nested within a 'host' study called 'STOP', which collaborated with different community and regional partners in many different ways during the planning and delivery of the intervention. In the majority of interventions, regardless of their target behavioural or health outcome, pharmacists received reimbursement for providing the intervention; this appears important in order for the intervention to be sustainable.^{32 36}

Organisation and delivery of the interventions

Sixteen interventions were delivered by the community pharmacy staff; one photoageing intervention³¹ was delivered by a research pharmacist employed by the

Table 1 Summary characteristics and outcomes of alcohol reduction interventions (further details are presented in online supplementary files 2–8)

| Study ID | Study characteristics | Description | Baseline behaviour | AUDIT total scores | | | FAST total scores | | | Global quality rating† | Effectiveness‡ | Cost-effectiveness | Differential effects§ |
|-----------------------------------|--|----------------------|-------------------------------|----------------------------|---------------|-----|----------------------------|------|----|------------------------|----------------|--------------------|-----------------------|
| | | | | Mean change from baseline* | 95% CI | N | Mean change from baseline* | SD | N | | | | |
| Dhital <i>et al</i> ²⁷ | Design: RCT | Brief alcohol advice | AUDIT Scores: 11.93 (SD 3.24) | −0.11 | −0.82 to 0.61 | 168 | | | | Strong | ↔ | NR | NR |
| | Duration: 12 weeks¶ Country: UK Number of pharmacies: 16 Number of participants: 407 Mean age: I:39.6; C: 40.5 % female: I: 47.8; C: 43.6 | Usual care control | AUDIT Scores: 11.53 (SD 3.19) | −0.74 p=0.24 | −1.47 to 0.00 | 158 | | | | | | | |
| Watson and Stewart ²⁸ | Design: RCT | Brief alcohol advice | FAST score ≥3: 29.2% | | | | 2.25 | 3.20 | 4M | Weak | ↔ | Cost analysis only | NR |
| | Duration: 26 weeks | Usual care | FAST score ≥3: 24.6% | | | | 0.50 | 0.71 | 2F | | | | |
| | Country: UK | | | | | | −1.25 | 2.87 | 4M | | | | |
| | Number of Pharmacies : 20 Number of participants: 69 Mean age: NR % female: I: 48.1; C: 57.1 | control | | | | | 0.75 NS | 1.67 | 8F | | | | |

*p Values were extracted directly from the study papers and relate to between group differences.

†Global rating: 'strong'=no 'weak' ratings, 'moderate'=one 'weak' rating and 'weak'=two or more 'weak' ratings.

‡Effectiveness was assessed using between group differences.

§Differential effects: age, gender, ethnicity or socioeconomic status (education, income, occupation, social class, deprivation or poverty).

¶From baseline to last follow-up.

↑, intervention effective; ↓, intervention not effective; ↔, no statistically significant between group difference; AUDIT, Alcohol Use Disorders Identification Test; C, control group; F, female; FAST, Fast Alcohol Screening Tool; I, intervention group; M, male; NR, not reported; NS, non-significant; RCT, randomised controlled trial.

Table 2 Summary characteristics and outcomes of smoking cessation interventions (further details are presented in online supplementary files 2–8)

| Study ID | Study characteristics | Description | Baseline behaviour | Quit rate* | Global quality rating† | Effectiveness‡ | Cost-effectiveness*** | Differential effects§ |
|-------------------------------------|---|---|---|---|------------------------|----------------|---------------------------------------|-----------------------|
| Bauld <i>et al</i> ²⁹ | Design: CBA Duration: 52 weeks¶ Country: UK Number of pharmacies: >200 Number of participants: 1785 Mean age: I: 44.0; C: 49.8 % female: I: 56.5; C: 65.5 | Individual pharmacy-based NHS smoking cessation service + NRT Group community-based NHS smoking cessation service + NRT | 21+ cigarettes/day: 396 (40.1%) 21+ cigarettes/day: 169 (41.6%) | 38/1374 (2.8%) 26/411 (6.3%) p=0.001 | Weak | ? | Yes both services compared to control | NR |
| Bock <i>et al</i> ³⁰ | Design: RCT Duration: 26 weeks Country: USA Number of pharmacies: 2 Number of participants: 299 Mean age: I1: 45.5; I2: 46.5; C: 42.3 % female: 59.0 | Smoking cessation training for pharmacists + tailored counselling using computer software + NRT Smoking cessation training for pharmacists + tailored counselling using computer software Observation only control (not randomised) | Number of cigarettes smoked/day: 18.2; Fagerström score: 5.3 Number of cigarettes smoked/day: 17.7; Fagerström score: 5.1 Number of cigarettes smoked/day: 13.8; Fagerström score: 4.9 | 28/100 (28.0%) 15/100 (15.0%) 8/99 (8.1%) p<0.01 | Moderate | ↑ | NR | NR |
| Burford <i>et al</i> ³¹ | Design: RCT Duration: 26 weeks Country: Australia Number of pharmacies: 8 Number of participants: 160 Mean age: I: 24.2; C: 25.1 % female: I: 68.7; C: 56.2 | Smoking cessation advice + computer-generated photoageing Smoking cessation advice | Fagerström score: 2.87; >21 cigarettes/day smoked: 10% Fagerström score: 2.96; >21 cigarettes/day smoked: 15% | 11/80 (13.8%) 1/80 (1.3%) p=0.003 | Moderate | ↑ | Yes | NR |
| Costello <i>et al</i> ³² | Design: RCT Duration: 5 weeks Country: Canada Number of pharmacies: 98 | 1 week then fortnightly visit for NRT plus 3 sessions brief behavioural counselling | HSI ≥3: 91.8% | 612/3503 (17.5%) | Weak | ↔ | NR | NR |

Continued

Table 2 Continued

| Study ID | Study characteristics | Description | Baseline behaviour | Quit rate* | Global quality rating† | Effectiveness‡ | Cost-effectiveness*** | Differential effects§ |
|--|---|--|--|--|------------------------|----------------|-----------------------|-----------------------|
| Crealey <i>et al</i> ¹⁹⁹⁰ <CE: Please check year is not matching with reference list.> | Number of participants:6987 Mean age:NR % female: I:54.4; C:54.9 | 5 weeks NRT at initial visit plus 1 session brief behavioural counselling | HSI ≥ 3: 91.4% | 604/3350 (18.0%) p=0.4 | | | | |
| | Design:nRCT Duration:26 weeks Country:UK Number of pharmacies:2 Number of participants:169 Mean age:NR % female: NR | Behavioural support, 67% (35/52) nicotine gum Nicotine gum only Control (expressed wish to stop smoking) | NR NR NR | 24/52 (46.2%) 3/48 (6.3%) 0/60 (0%) p<0.01 (I vs C) | Weak | ↑ | Yes | NR |
| | | | | | | | | |
| Hoving <i>et al</i> ³⁴ | Design: RCT Duration:52 weeks Country:Netherlands Number of pharmacies:65 Number of participants:545 Mean age:I:46; C:47 % female: I:53; C:54 | Computer-generated tailored advice 'Thank you' letter control | Number of cigarettes smoked/day: 22 Number of cigarettes smoked/day: 21 | 2/256 (0.8%) 2/289 (0.7%) NS | Strong | ↔ | NR | NR |
| | | | | | | | | |
| Howard-Pitney <i>et al</i> ³⁵ | Design: RCT Duration:26 weeks Country:USA Number of pharmacies:5 Number of participants:410 Mean age:I:36.3; C:34.7 % female: I:1; C:1 | Advice and support + nicotine patch Advice and support + placebo patch | Number of cans chewed/week: 3.9 Number of cans chewed/week: 4.1 | 78/206 (37.9%) 69/204 (33.8%) p<0.40 | Moderate | ↔ | NR | NR |
| Maguire <i>et al</i> ³⁶ | Design: RCT Duration:52 weeks Country:UK Number of pharmacies:51 Number of participants:484 Mean age:I:42; C:38 % female: I:40; C:44 | Behavioural support, 87% (230/265) NRT Ad hoc advice, 84% (183/219) NRT | Number of participants 10–20 cigarettes/day: 197/265 Number of participants 10–20 cigarettes/day: 121/219 | 38/265 (14.3%) 6/219 (2.7%) p < 0.001 | Weak | ↑ | NR | NR |
| | | | | | | | | |

Continued

Table 2 Continued

| Study ID | Study characteristics | Description | Baseline behaviour | Quit rate* | Global quality rating† | Effectiveness‡ | Cost-effectiveness*** | Differential effects§ |
|---------------------------------------|---|--|--|--|------------------------|---------------------|-----------------------|-----------------------|
| Mochizuki <i>et al</i> ³⁷ | Design: RCT Duration:12 weeks Country:Japan Number of pharmacies:14 Number of participants:28 Mean age:I:44.1; C:49.1 % female: I:18.2; C:18.8 | Nicotine gum plus advice on usage, initial and follow-up cessation advice Nicotine gum plus advice on usage | Number of cigarettes smoked/day: 23.0; Fagerström score: 4.56 Number of cigarettes smoked/day: 25.7; Fagerström score: 6.31 | 5/11 (45.5%) 5/16 (31.3%) OR=1.83, NS | Strong | ↔ | NR | NR |
| Sinclair <i>et al</i> ³⁸ | Design: RCT Duration:36 weeks Country:UK Number of pharmacies:62 Number of participants:492 Mean age:I:41.7; C:41.5 % female: I:61.2; C:62.7 | Training pharmacists/ assistants in smoking cessation behaviour change + NRT Standard professional pharmacy support + NRT | Fagerström score: 5.2 Fagerström score: 5.2 | 26/217 (12.0%) 19/257 (7.4%) p=0.089 | Strong | ↔ | Yes | NR |
| Sonderskov <i>et al</i> ⁴⁰ | Design: RCT Duration:26 weeks Country:Denmark Number of pharmacies:42 Number of participants:522 Mean age:I(21 mg):39.1; C(21 mg):39.9; I(14 mg):38.2; C(14 mg):38.9 % female: I(21 mg):47.5; C(21 mg):52.5; I(14 mg):51.7; C(14 mg):48.3 | 21 mg nicotine patches Placebo 14 mg nicotine patches Placebo | Fagerström score: 7.0 Fagerström score: 8.1 Fagerström score: 6.1 Fagerström score: 6.1 | 15/132 (11.4%) 6/142 (4.2%) p<0.05 27/119 (22.7%) 23/125 (18.4%) NS | Strong | ↑ 21 mg; ↔ 14 mg | NR | No (gender) |

Continued

Table 2 Continued

| Study ID | Study characteristics | Description | Baseline behaviour | Quit rate* | Global quality rating† | Effectiveness‡ | Cost-effectiveness*** | Differential effects§ |
|---------------------------------|--|---|--|---|------------------------|----------------|-----------------------|-----------------------|
| Vial <i>et al</i> ³⁹ | Design: RCT Duration: 52 weeks Country: Australia Number of pharmacies: 9 Number of participants: 102 Mean age: 51.0 % female: 11.41; 12.54; C:36 | Pharmacy-based nicotine patches plus weekly counselling Hospital outpatient clinic nicotine patches plus weekly counselling Minimal intervention (written and verbal information at baseline) | Fagerström score: 5.79 Fagerström score: 5.94 Fagerström score: 6.33 | 4/21 (19.0%) 5/21 (23.8%) 1/22 (4.5%) NS | Weak | ↔ | NR | NR |

*p Values were extracted directly from the study papers and relate to between group difference.

†Global rating: 'strong'=no 'weak' ratings, 'moderate'=one 'weak' rating and 'weak'=two or more 'weak' ratings.

‡Effectiveness was assessed using between group differences.

§Differential effects: age, gender, ethnicity or socioeconomic status (education, income, occupation, social class, deprivation or poverty).

¶From baseline to last follow-up.

?, Unable to assess effectiveness/cost-effectiveness; †, intervention effective; ‡, intervention not effective; ↔, no statistically significant between group difference; C, control group; CBA, controlled before-after study; Fagerström score, 0–10, higher score=greater nicotine dependence; HHS, Heaviness of Smoking Index, higher score indicates greater number of cigarettes smoked per day and smoking first cigarette within 5 min of waking; I, intervention group; NHS, National Health Service; NR, not reported; nRCT, non-randomised controlled trial; NRT, nicotine replacement therapy; NS, non-significant; RCT, randomised controlled trial.

local university in collaboration with the community pharmacist, who delivered standard smoking cessation advice (see online supplementary file 5). Another smoking cessation intervention was conducted by a research pharmacist as part of an MSc project; the research pharmacist delivered the hospital-based intervention programme, and the community pharmacists delivered the community-based intervention programme.³⁹ One smoking cessation intervention involved the postal delivery of a computer-generated letter.³⁴ Most studies included standardised staff training, although this was usually brief (ranging from 2 h to 2 days). Two smoking cessation studies mentioned they also included role play as part of the training,^{30 35} and two weight management studies reported 'practical tasks' as part of the training.^{43 45}

In terms of quality assurance, one alcohol reduction intervention provided a 2 h evening follow-up training session during the intervention to address challenges and share learning across the pharmacists who were delivering the intervention.²⁷ In two smoking cessation studies,^{33 36} a researcher visited the pharmacists after the group training session, to provide support and to address any queries they had in implementing the training. In one smoking cessation study that was organised by a pharmaceutical company,⁴⁰ the company contacted pharmacies at least once a week during the intervention.

Behaviour change strategies used in the interventions

Seven studies reported that a behavioural theory/model informed the intervention and provided details of behaviour change strategies used; six studies only reported details of behaviour change strategies used; six studies reported no relevant information (see online supplementary file 6). The most commonly reported theoretical model was the Transtheoretical ('Stages of Change') Model, which was reported by six studies; motivational interviewing was reported by five studies. The descriptions available did not allow for the coding of specific aspects of theory and behavioural content. Using the Behaviour Change Wheel,¹⁸ the intervention functions of the majority of interventions were 'education' and 'enablement'. In addition, interventions that included the provision of NRT or commercial weight management programmes or products free of charge, were also deemed to include 'incentivisation'. Using the policy category of the Behaviour Change Wheel,¹⁸ all the interventions were categorised as 'service provision'. Six of these interventions also included 'communication/marketing'. No other policy categories were identified. Using the Nuffield intervention ladder,¹⁹ most interventions were coded as 'enable choice'.

EFFECTS OF INTERVENTIONS

Alcohol reduction interventions (n=2)

There were two RCTs of brief alcohol reduction interventions (table 1) compared with usual care or

**Table 3** Summary characteristics and outcomes of weight management interventions (further details are presented in online supplementary files 2–8)

| Study ID | Study characteristics | N | Description | BMI (kg/m ²) | | WC (cm) | | WT (kg) | | Global quality rating* | Effectiveness† | Cost-effectiveness | Differential effects‡ |
|---|---|-----|---|----------------------------|--------------|----------------------------|-----|----------------------------|--------------|------------------------|----------------|---|---|
| | | | | Mean change from baseline§ | SD/95% CI | Mean change from baseline§ | SD | Mean change from baseline§ | SD/95% CI | | | | |
| Ahrens <i>et al</i> ⁴¹ 2011 | Design: RCT | 45 | Meal replacement diet | NR | NR | −8.08 | NR | −5.6 | NR | Weak | ↔ | NR | NR |
| | Duration: 22 weeks¶ Country: USA Number of pharmacies: 1 Number of participants: 95 Mean age: I: 47.6; C: 47.8 % female: 87 Baseline BMI: I: 29.5; C: 29.0 | 43 | Low calorie diet | NR | NR | −7.82 | NR | −5.2 | NR | | | | |
| Bush <i>et al</i> ⁴² | Design: CBA | 60 | Pharmacy-based diet + physical activity | −1.3 | 0.4 | −6.5 | 1.6 | −3.4 | 1.1 | Weak | ? | Unclear which service was more cost effective | Yes, demographics of participants differed significantly between settings |
| | Duration: 15 weeks¶ Country: UK Number of pharmacies: 12 Number of participants: 451 Mean age: I: 38.9; C: 42.6 % female: I: 87; C: 85 Baseline BMI: I: 33.0; C: 35.6 | 22 | GP-based diet + physical activity | −0.8 | 0.7 | −4.9 | 2.6 | −2.3 | 1.9 | | | | |
| Jolly <i>et al</i> ^{43**} | Design: RCT | 70 | Pharmacy-based diet + physical activity | −0.31 | −0.7 to 0.0 | NR | NR | −0.66 | −1.7 to 0.4 | Moderate | ↔ | Cost analysis only, commercial organisations lower cost than GP and pharmacy-based services | No (gender) |
| | Duration: 52 weeks¶ Country: UK | | | | | | | | | | | | |
| | Number of pharmacies:NR | 100 | Exercise only control | −0.45 | −0.8 to −0.1 | NR | NR | −1.08 | −2.1 to −0.1 | | ‡ | | |
| | Number of participants: 740 | 100 | Weight Watchers | −1.17 | −1.7 to −0.7 | NR | NR | −3.46 | −4.8 to −2.1 | | ↑ | | |
| | Mean age: Ph: 48.9; Ex: 49.7; WW: 50.7; SW: 48.8; RC: 48.8; NHS SD: 48.8; GP: 50.5; POC: 47.5 | 100 | Slimming World | −0.71 | −1.0 to −0.4 | NR | NR | −1.89 | −2.9 to −0.9 | | ↔ | | |
| | % female: Ph: 73; Ex: 75; WW: 72; | 100 | Rosemary Conley | −0.75 | −1.1 to −0.3 | NR | NR | −2.12 | −3.4 to −0.9 | | ↔ | | |
| | | 100 | NHS Size Down | −0.67 | −1.0 to −0.3 | NR | NR | −2.45 | −3.6 to −1.3 | | ↔ | | |
| | | 70 | GP | −0.32 | −0.7 to 0.1 | NR | NR | −0.83 | −2.0 to 0.4 | | ↔ | | |
| | | 100 | Participants own choice | −0.90 | −1.3 to −0.5 | NR | NR | −2.15 | −3.4 to −0.9 | | ↔ | | |

Continued

Table 3 Continued

| Study ID | Study characteristics | N | Description | BMI (kg/m ²) | | WC (cm) | | WT (kg) | | Global quality rating* | Effectiveness† | Cost-effectiveness | Differential effects‡ |
|--------------------------------------|--|----|----------------------------------|----------------------------|-----------|----------------------------|------|----------------------------|-----------|------------------------|----------------|--------------------|-----------------------|
| | | | | Mean change from baseline§ | SD/95% CI | Mean change from baseline§ | SD | Mean change from baseline§ | SD/95% CI | | | | |
| Malone and Alger-Mayer ⁴⁴ | SW: 65; RC: 69; NHS SD: 64; GP: 67; POC: 70 Baseline BMI: P: 33.4; Ex: 33.9; WW: 34.0; SW: 33.8; RC: 33.4; NHS SD: 33.8; GP: 33.1; POC: 33.4 Design: nRCT Duration: 26 weeks¶ Country: USA Number of pharmacies: NR Number of participants: 30 Mean age: I: 44.9; C: 42.8 % female: I: 93; C: 80 Baseline BMI: I: 48.3; C: 42.8 | 15 | Pharmacist support + orlistat | NR | NR | NR | NR | -3.5 | 2.9 | Weak | ↔ | NR | NR |
| | orlistat + usual outpatient care | 15 | orlistat + usual outpatient care | NR | NR | NR | NR | -3.0 | 5.2 | | | | |
| Phimarn et al ⁴⁵ | Design: RCT Duration: 16 weeks¶ Country: Thailand Number of pharmacies: 1 Number of participants: 66 Mean age: I: 60.1; C: 59.1 % female: I: 75.8; C: 84.8 Baseline BMI: I: 27.5; C: 27.7 | 33 | Pharmacist individual support | -0.8 | 0.07 | 0.1 | 0.03 | -0.82 | 0.29 | Strong | ↔ | NR | NR |
| | | 33 | Primary care unit group support | 0.19 | 0.04 | -0.28 | 0.08 | 0.92 | 0.19 | | | | |

*Global rating: 'strong'=no 'weak' ratings, 'moderate'=one 'weak' rating and 'weak'=two or more 'weak' ratings.

†Effectiveness was assessed using between group differences.

‡Differential effects: age, gender, ethnicity or socioeconomic status (education, income, occupation, social class, deprivation or poverty).

**All intervention groups in the Jolly trial were compared to the exercise only control group (intervention groups were not directly compared).

§p Values were extracted directly from the study papers and relate to between group differences.

¶From baseline to last follow-up.

↓, intervention not effective; ↑, intervention effective; ↔, no statistically significant between group difference; ?, unable to assess effectiveness/cost-effectiveness; BMI, body mass index; C, control group; CBA, controlled before-after study; Ex, exercise only control; GP, general practitioner; NHS SD, NHS Size Down; NHS, National Health Service; NR, not reported; nRCT, non-randomised controlled trial; NS, non-significant; Ph, Pharmacy-based diet + physical activity; POC, participants own choice; RC, Rosemary Conley; RCT, randomised controlled trial; SW, Slimming World; WC, waist circumference; WT, weight; WW, Weight Watchers.

leaflet-only control (see online supplementary files 7 and 8). One RCT²⁷ used the Alcohol Use Disorders Identification Test total scores (AUDIT), and reported a baseline AUDIT score of 11.93. The other RCT²⁸ used the Fast Alcohol Screening Tool (FAST), and reported 29.2% of participants scoring ≥ 3 at baseline. Possible 'harmful or hazardous' alcohol consumption, but not alcohol dependence is indicated with an AUDIT score 8–19 or a FAST score of 3–16. Global quality ratings were 'strong' for one study²⁷ and 'weak' for the other,²⁸ which was a small pilot study. Both studies involved one-to-one contact with the pharmacist. Dhital *et al*²⁷ encouraged self-directed behaviour change; the intervention included reflection and feedback of the AUDIT score.

Behavioural outcomes

Neither intervention significantly reduced alcohol scores compared with control. At 12 weeks, the AUDIT total change score did not differ significantly between the two groups and did not change significantly between baseline and follow-up in either group. Twelve-week AUDIT between group difference, adjusted for pharmacist gender, age, ethnicity and education, was -0.57 (95% CI -1.59 to 0.45). There was no significant difference between FAST score for the intervention group compared with control at 3 or 6 months, adjusted for baseline FAST: the difference between groups was -1.84 (95% CI -4.49 to 0.82). At 6 months, there was substantially lower follow-up of intervention participants (22.2%) compared with control participants (33.3%).

Costs

Cost-effectiveness of community pharmacy-based brief alcohol reduction interventions cannot be ascertained; only one pilot study²⁸ reported direct intervention costs.

Differential effects by demographic or socioeconomic factors

One pilot study²⁸ reported change in FAST scores by gender within the intervention and control groups. However, the study was not powered to detect differences between the two groups.

Smoking cessation interventions (n=12)

There were 10 RCTs,^{30–32 34–40} 1 nRCT³³ and 1 CBA²⁹ of smoking cessation interventions (table 2). Global quality ratings were 'strong' for four studies, 'moderate' for three studies, and 'weak' for five studies. Eleven studies carried out analyses with the assumption that those lost to follow-up had not stopped smoking. Half (6/12) the smoking cessation interventions relied on self-reported change in smoking behaviours,^{32 34 37 38–40} and half used biochemical measures (carbon monoxide (CO) or cotinine levels).^{29–31 33 35 36}

Ten studies included NRT (in either the intervention or control group or both).^{29 30 32 33 35–40} Seven studies evaluated some form of behavioural support.^{30 32–34 36–38} Two studies evaluated the effect of intervention setting;

one study assessed behavioural support plus NRT provided in a hospital outpatient setting compared with pharmacy setting.³⁹ Another compared individual pharmacy-based behavioural support plus NRT with group-support provided in a community setting.²⁹ One study³¹ evaluated the effect of a photoageing intervention.

Despite a variety of different components being evaluated within the individual interventions, the studies were grouped together to assess the effectiveness of any type of community pharmacy-delivered intervention for smoking cessation compared with either an active control or a non-active/usual care comparator. 'Usual care' varied between studies but was, in general, a minimal intervention, such as observation only, ad hoc smoking cessation advice or a thank you letter. However, in one study, the control group received placebo nicotine patches⁴⁰ and in two studies the control group received standard cessation advice plus NRT.^{36 38}

Behavioural outcomes

Five of the 12 studies demonstrated effectiveness compared with control. In addition, Bauld *et al*²⁹ evaluated one-to-one pharmacist support with group-based smoking cessation clinics based in the community; the group-based service attracted fewer clients but was more effective.

The five effective studies included:

- ▶ An American RCT³⁰ of additional training to pharmacists to enable them to provide a tailored counselling service with and without NRT, compared with a non-randomised control group that received observation only, showed a significant increase in validated 7-day point prevalence at 6 months (28% for counselling and NRT, 15% for counselling, 8% for control).
- ▶ An Australian RCT³¹ of a computer-generated photoageing service (demonstrating the detrimental effects on facial physical appearance of smoking) in addition to standard smoking cessation advice from a pharmacist, was effective in stopping young people (mean age 24 years) smoking compared to control using CO-validated measures (13.8% $n=22/80$ vs 1.3% $n=11/80$) at 6 months.
- ▶ A cost-effectiveness study³³ in two UK pharmacies compared a behavioural intervention group based on the Pharmacist Action on Smoking (PAS) model with a control group that received nicotine gum, and another control group who expressed a wish to stop smoking. At 6 months, there was a statistically significant difference in cessation rates between intervention and control groups. Six-month CO-verified abstinence was 46% in the intervention group, 6% in the nicotine gum control group, and 0% in the control group that expressed a wish to stop smoking.
- ▶ A UK RCT³⁶ compared an intervention based on the PAS model to ad hoc smoking cessation advice; over 80% in each group also had NRT. The PAS intervention significantly increased validated smoking

cessation compared with control at 12 months (14.3% vs 2.7%).

- A Danish study⁴⁰ evaluated the effect of two different strengths of nicotine patches compared to placebo. Those smoking ≥ 20 /day at baseline were randomised to 21 mg patches or placebo, those smoking < 20 /day at baseline were randomised to 14 mg patches or placebo. Self-reported point prevalence included participants who had one episode of smoking (< 6 days). At 26 weeks, the intervention was effective for those smoking ≥ 20 /day at baseline (11% vs 4.2%) but not effective for lighter smokers (22.7% vs 18.4%) compared with the respective placebo groups.

Meta-regression and meta-analysis

Meta-regression of ORs of smoking cessation between the intervention and the control groups was undertaken; in model 1, a random effects model was fitted including all the RCTs. The pooled OR for the intervention effects was 1.85 (95% CI 1.25 to 2.75), an indication of the positive effect of the interventions on smoking cessation. However, there was 72% unexplained differences between the studies. In model 2, a meta-regression model was fitted accounting for whether a study had an active comparator or non-active/usual care comparator. The pooled ORs were 1.21 (95% CI 0.86 to 1.71) and 2.56 (95% CI 1.45 to 4.53) for the active comparator or non-active/usual care comparator, respectively (figure 2).

As expected, there was a larger effect when compared with non-active/usual care comparator than with active comparator. The proportion of unexplained heterogeneity reduced to 52%. In model 3, a meta-regression model was fitted accounting for whether a study had an

active comparator or a non-active/usual care comparator, and also the intervention duration; the unexplained heterogeneity reduced to 27.2% with a non-significant Q-statistic test (10.99, $p < 0.2026$). In model 4, quality rating was accounted for; quality rating did not appear to contribute much to the model after accounting for intervention duration, and whether a study had an active comparator or a non-active/usual care comparator. Figure 3 shows a meta-analysis of smoking cessation accounting for global quality rating, and shows that most variations between studies are from studies rated as 'moderate' or 'weak' quality. A funnel plot demonstrated asymmetry, with larger studies showing effects closer to the null than smaller studies. Such a pattern is compatible with publication bias, on the assumption that smaller studies with uninteresting effects are withheld from publication. However, the funnel plot must be interpreted with caution, taking into account that it contains only 10 studies, which is the recommended study size threshold for creating such plots.¹¹

Costs

Four studies reported cost-effectiveness analyses; the costs and benefits differed between the studies, and costs years ranged from 1995 to 2011, making comparisons across the analyses difficult. All four studies used quit rates observed within the trials, these ranged from 2.8% to 12% for UK pharmacist-based behavioural support with NRT.

By comparison with a self-quit attempt, the incremental cost per Quality Adjusted Life Year was £2600 for pharmacy one-to-one counselling, and £4800 for group community-based NHS smoking cessation service.²⁹ Incremental Cost Effectiveness Ratios (ICER) per




Figure 2 Meta-analysis of smoking cessation accounting for whether active comparator or non-active comparator.




Figure 3 Meta-analysis of smoking cessation accounting for global quality rating.

additional quitter ranged from £79 to £509 for pharmacist-based behavioural support with NRT. The ICER per additional quitter using photoageing was \$A46 (Australian dollars 2011).³¹ In summary, three UK pharmacy-delivered interventions appeared cost effective across a range of quit rates, and an Australian photoageing intervention was cost effective compared to standard advice among young adults.

Differential effects by demographic or socioeconomic factors

A Danish study⁴⁰ evaluated the effect of two different strengths of nicotine patches compared with placebo. There were no differences in smoking cessation rates between men and women according to starting dose and treatment.

Weight management interventions (n=5)

There were three RCTs,^{41 43 45} one nRCT⁴⁴ and one CBA⁴² of weight management interventions (table 3). Global ratings were 'strong' for one study, 'moderate' for one study, and 'weak' for three studies. Three studies^{42 43 45} compared a pharmacy-based intervention with similar interventions in other primary care settings, and commercial programmes in community settings. One study⁴¹ compared a meal replacement diet with a conventional low-energy diet (identical recommended total daily energy intake); both interventions were set in a pharmacy. One small study⁴⁴ assessed the added value of community pharmacy support for an obesity

management intervention that included orlistat and an outpatient nutrition programme.

Anthropometric outcomes

Three studies reported body mass index (BMI), three studies reported waist circumference (WC) and all five studies reported weight (WT). None of the studies found a significant difference in favour of a pharmacy-delivered intervention compared with the comparator, for any anthropometric outcome. However, all comparators are 'active' interventions (smoking cessation studies demonstrated larger effect when compared with non-active controls compared to active controls). One UK RCT⁴³ compared seven groups (Weight Watchers, Slimming World, Rosemary Conley, Size Down an NHS community-based group, GP, Pharmacy, participants' own choice to an exercise-only control group). This study compared each intervention group with a control group, and was not designed to directly compare the active interventions which were delivered across different settings. All, except the GP and pharmacy groups, resulted in significant weight loss at 1 year compared with baseline. Mean weight loss at 1 year, with baseline value used for imputation, was 0.8 kg (SD 4.7 kg) for primary care (GP and pharmacy) and 2.5 kg (SD 6.2 kg) for commercial programmes. Only the Weight Watchers group demonstrated significant weight loss at 1 year compared to control.

One CBA⁴² study compared diet and physical activity in a pharmacy to a GP-based intervention: both groups

appeared to reduce BMI, WC and WT at follow-up (statistical significance not reported). Despite participants choosing the service, there was very high attrition (93%). One study⁴¹ demonstrated significant and similar amounts of weight loss from baseline to follow-up for participants in a meal replacement group, or a low-calorie diet group (both pharmacy-delivered). In another study, pharmacy-based support in addition to orlistat did not improve weight loss.⁴⁴ Another study demonstrated no significant improvement in weight from baseline to follow-up for participants receiving group-based support in a primary care unit compared with individual support from a pharmacist.⁴⁵

Costs

Two trials reported intervention costs, one of which also reported costs per kg weight lost.⁴² The Jolly *et al*⁴³ trial reported similar costs (£112) for both the pharmacy group and the GP group; both settings had higher costs compared with commercial weight management programmes (£71–£77), the NHS community-based group costs fell in-between at £92.

A study⁴² of weight management programmes based in pharmacy or GP settings reported costs ((£126.90 per participant (n=183) in the pharmacy intervention and £100.60 per participant (n=268) in the GP intervention)), that were broadly similar to that of the pharmacy-based group in the Jolly trial. It is unclear which provider type delivered the intervention more cost-effectively; at session 12, the ICER (£ per kg per participant) cost –£8.29 through pharmacy providers (favours GP). Conversely, at the final session 15, the ICER was £2.91 through GP providers (favours Pharmacy).

Differential effects by demographic or socioeconomic factors

In a study of weight management programmes in various commercial, primary care and NHS settings, there was no statistically significant interaction between gender and the type of weight management programme.⁴³ Bush *et al*⁴² compared a weight management programme set in pharmacies with the same programme set in GP surgeries. Female participants in GP surgeries lost a significantly larger proportion of their initial weight than female participants in pharmacies; participants aged 40–49 years lost a greater proportion of their initial weight at GP providers than at pharmacy providers.

DISCUSSION

Community pharmacy-delivered smoking cessation interventions including behavioural support and/or NRT, are effective and cost effective, particularly when compared with usual care. The pooled ORs for smoking cessation were 1.21 (95% CI 0.86 to 1.71) and 2.56 (95% CI 1.45 to 4.53) for active control and usual care, respectively. The heterogeneity of types of interventions precluded the ability to evaluate effectiveness by specific types of

interventions. There was little evidence comparing pharmacy-delivered smoking cessation with smoking cessation delivered in other settings. This was in contrast with the majority of the weight management evidence which compared active interventions in various settings. Although there was no significant difference in weight loss between active interventions in different settings; pharmacy-based interventions produced similar amounts of weight loss (3–5 kg) from baseline to short-term follow-up (6 months or less), compared with active interventions in other primary care settings. There was insufficient evidence to evaluate community pharmacy-delivered interventions for alcohol reduction.

Community pharmacy-delivered smoking cessation interventions are cost-effective (compared to self-quit or standard care) across a range of quit rates. Cost-effectiveness of pharmacy-delivered weight management interventions is unclear; they have similar provider costs to those delivered in other primary care settings, which are greater than those delivered by commercial organisations. This review aimed to extract information on intervention costs and potential cost savings; however, it is not a review of economic evaluations and, as such, the methods of the economic evaluations are not critically appraised; we simply report the results of the economic evaluations that were conducted alongside included interventions.

Evidence suggests that duration of intervention is a predictor of effectiveness, for both the smoking cessation and weight loss studies; in the case of weight management, longer term weight loss may differ by setting. Regression analysis showed that duration of intervention accounted for some heterogeneity across the smoking cessation studies. One longer term weight management study compared interventions in a range of primary care and commercial settings; all except the pharmacy and GP groups resulted in significant weight loss at 1 year compared with baseline. The data reported in the studies identified for this review highlight the potential importance of predictors of success.

In terms of the effects of the interventions on health inequalities, some studies examined demographic and/or socioeconomic factors at recruitment stage, as potential predictors of outcomes within group, and/or to explain differences in retention. However, none of the studies reported subgroup analysis of treatment effect by SES. Three studies adopted a targeted approach to addressing inequality, by recruiting participants from deprived areas, and compared a pharmacy setting with other settings. In two of these 'targeted' studies (one smoking, one weight) the participants self-selected the service; there were demographic and socioeconomic differences between participants who self-selected treatment by setting. The evidence shows that the community pharmacy is an appropriate and feasible setting to deliver a range of public health interventions, and this setting has the potential to reach those most in need.

The original analysis plan included an examination of any potential relationships between intervention effectiveness and behaviour change strategies and/or models used, also whether any patterns existed between effective interventions and implementation factors (eg, pharmacist training or resource intensity) or the organisation and delivery of service (eg, stakeholder involvement). Unfortunately, the lack of relevant data reported meant that this analysis could not be undertaken. It is worth noting that the majority of interventions were implemented within the political context of extending the public health role of pharmacists.

These findings build on previous work; looking to the future, there is a Cochrane review⁽⁴⁶⁾ in progress with a broader remit than this review; it evaluates the effectiveness of a wider variety of health promotion, or health behaviour interventions, set in community pharmacy. This work will further develop the evidence base; in particular, evidence regarding implementation, organisation and delivery of other types of public health interventions which may be transferable to alcohol reduction, smoking cessation and weight management interventions.

Implications for policy and practice. The evidence shows a range of types of smoking cessation interventions that are feasible and effective within community pharmacies, and supports the commissioning of smoking cessation services in a community pharmacy setting. Smoking cessation services, contracted as a core part of the national contract, or part of a national 'advanced' service, may well be a reasonable option. In addition, the evidence shows that weight management services are no less effective compared with those delivered in other primary care settings. Therefore, given the potential reach, effectiveness and associated costs of these interventions, commissioners may consider using community pharmacies to help deliver some of their smoking cessation and weight management services.

Implications for future research: Further research is required to assess the effectiveness of community pharmacy-delivered alcohol reduction interventions, and more research is needed on the cost-effectiveness of community pharmacy-delivered alcohol, smoking and weight management interventions compared with other providers. It appears that the duration of intervention impacts on effectiveness, and this is likely to impact on cost-effectiveness. There is a lack of evidence regarding the effect of community pharmacy-based interventions for alcohol reduction, smoking cessation and weight loss on health inequalities. Targeted intervention studies provided some evidence that adults accessing pharmacies are a distinct group that may not access other primary care or commercial organisations. This evidence is derived from participants who self-selected the intervention and setting. However, more research is required on the reach of public health interventions delivered from a community pharmacy setting. Future studies should be sufficiently powered to detect small changes in behavioural and health outcomes and measure the equity

effects of these small changes at a population level. Future studies should assess and report sociodemographic and socioeconomic variables, behaviour change strategies and models, implementation factors, the organisation and delivery of interventions, and costs.

CONCLUSIONS

The evidence demonstrates that the community pharmacy is an appropriate and feasible setting to deliver a range of public health interventions. Community pharmacy-delivered smoking cessation interventions are effective and cost effective, particularly when compared with usual care. Evidence from a heterogeneous group of weight management interventions suggest that community pharmacy-delivered weight management interventions are as effective as similar interventions in other primary care settings, at least in the short term, and have similar provider costs. There is insufficient evidence to assess the effectiveness of community pharmacy-based interventions for alcohol reduction. The impact of community pharmacy-delivered interventions on inequalities in priority public health conditions is unclear.

Author affiliations

¹School of Medicine, Pharmacy and Health, Durham University Queen's Campus, Stockton-on-Tees, UK

²FUSE, UKCRC Centre for Translational Research in Public Health, Newcastle University, Newcastle Upon Tyne, UK

³Wolfson Research Institute for Health and Wellbeing, Durham University Queen's Campus, Stockton-on-Tees, UK

⁴Centre for Health and Inequalities Research, Department of Geography, Durham University, Durham, UK

⁵Institute of Health & Society, Newcastle University, Newcastle Upon Tyne, UK

⁶Blizard Institute, Barts and The London School of Medicine and Dentistry, London, UK

⁷Sheffield Business School, Sheffield Hallam University, Sheffield, UK

Twitter Follow Adam Todd at @adam.todd138

Acknowledgements The authors acknowledge the contribution of the members of Review Advisory group for their advice regarding the protocol for the review and study identification: Claire Anderson (Professor of Social Pharmacy, Head of Division of Social Research in Medicines and Health, University of Nottingham), Christine Bond (Professor, Chair in General Practice and Primary Care, University of Aberdeen), Mark Burdon (Community Pharmacist, member of Pharmaceutical Services Negotiating Committee), Carol Hall (retired, patient, member of the pharmacy-specific patient and public involvement group), Eileen Kaner (Professor and Director of the Institute of Health & Society, expertise in alcohol misuse, University of Newcastle), Elena Ratschen (Lecturer in Epidemiology/Tobacco Control, University of Nottingham), Pat Simpson (retired, patient, member of the pharmacy-specific patient and public involvement group) and Margaret Watson (Senior Research Fellow, Pharmacy, University of Aberdeen). The coauthors acknowledge the contribution of Julian Higgins (Professor of Evidence Synthesis, University of Bristol) for advice on evidence synthesis. The coauthors also acknowledge the contribution of Nasima Akhter (Post-Doctorate, Durham University) for statistical support.

Contributors TJB was responsible for data collection, and contributed to analysis, synthesis and interpretation. She led the drafting of this article. CDS was responsible for overall design, coordination and project management. She provided methodological and conceptual direction, and contributed to analysis, synthesis and interpretation. AT provided methodological, analysis

and synthesis input, and contributed to data collection and synthesis. AH, CB, FFS and LS provided methodological, analysis and synthesis input. AK contributed to analysis. HJM designed and conducted the searches with support from LS, and contributed to data collection. CO contributed to data collection and synthesis. SS and LN contributed to data collection. All authors contributed to revised successive drafts of this article and approved the final version submitted for publication.

Funding This project was funded by the National Institute for Health Research Public Health Research Programme (project number 12/153/52): <http://www.nets.nihr.ac.uk/projects/phr/1215352>.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- World Health Organization. *Global strategy on diet, physical activity and health*. Geneva: World Health Organization, 2004.
- Bambra C, Joyce K, Maryon-Davies A. Strategic Review of Health Inequalities in England post-2010 London; (Marmot review) (Task Group 8: Priority Public Health Conditions: Final Report). London: University College London Institute of Health Equity, 2009.
- Todd A, Copeland A, Husband A, *et al*. Access all areas? An area-level analysis of accessibility to general practice and community pharmacy services in England by urbanity and social deprivation. *BMJ Open* 2015;5:e007328.
- Todd A, Copeland A, Husband A, *et al*. The positive pharmacy care law: an area-level analysis of the relationship between community pharmacy distribution, urbanity and social deprivation in England. *BMJ Open* 2014;4:e005764.
- Pharmaceutical Services Negotiating Committee. Service specifications and resources. <http://psnc.org.uk/service-search-results/?location-of-service=293&type-of-service=387&method-of-commissioning=&funding-source=&wqsfsubmit=Filter> (accessed 17 Jun 2015).
- Peterborough Pharmaceutical Needs Assessment. Peterborough City Council, 2015. <https://www.peterborough.gov.uk/upload/www.peterborough.gov.uk/healthcare/public-health/Healthcare-PublicHealth-PeterboroughPNA2015.pdf?inline=true> (accessed 30 Jun 2015).
- Department of Health. *Pharmacy in England: building on strengths delivering the future*. London: Department of Health, 2008.
- Gordon J, Watson M, Avenell A. Lightening the load? A systematic review of community pharmacy-based weight management interventions. *Obes Rev* 2011;12:897–911.
- Watson MC, Blenkinsopp A. The feasibility of providing community pharmacy based services for alcohol misuse: a literature review. *Int J Pharm Pract* 2009;17:199–205.
- Sinclair HK, Bond CM, Stead LF. Community pharmacy personnel interventions for smoking cessation. *Cochrane Database Syst Rev* 2004;(1):CD003698.
- Higgins JPT, Green S. *Cochrane handbook for systematic reviews of interventions version 5.1.0*. The Cochrane Collaboration, 2011.
- Todd A, Moore HJ, Husband AK, *et al*. Community pharmacy interventions for public health priorities: protocol for a systematic review of community pharmacy delivered smoking, alcohol and weight management interventions. *Syst Rev* 2014;3:93.
- Moher D, Liberati A, Tetzlaff J, *et al*. The PRISMA Group. Preferred reporting items for systematic reviews and meta-Analyses: the PRISMA Statement. *BMJ* 2009;339:b2535.
- Welch V, Petticrew M, Tugwell P, *et al*. PRISMA-equity 2012 extension: reporting guidelines for systematic reviews with a focus on health equity. *PLoS Med* 2012;9:e1001333.
- Cochrane Effective Practice and Organisation of Care (EPOC). *What study designs should be included in an EPOC review? EPOC Resources for review authors*. Oslo: Norwegian Knowledge Centre for the Health Services, 2013. <http://epoc.cochrane.org/epoc-specific-resources-review-authors> (accessed 30 Jun /2015).
- Egan M, Bambra C, Petticrew M, *et al*. Reviewing evidence on complex social interventions: appraising implementation in systematic reviews of the health effects of organisational-level workplace interventions. *J Epidemiol Community Health* 2009;63:4–11.
- Bambra CL, Hillier-Brown FC, Cairns JM, *et al*. How effective are interventions at reducing socioeconomic inequalities in obesity among children and adults? Two systematic reviews. *Public Health Res* 2015;3:1–446.
- Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 2011;6:42.
- Policy Process and Practice. *Public Health: ethical issues*. London: Nuffield Council on Bioethics, 2007.
- Effective Public Health Practice Project. Effective public health practice project quality assessment tool for quantitative studies. <http://www.epphp.ca/tools.html> (accessed 30 Jun 2015).
- Cochrane Public Health Group. Guidance for authors. <http://ph.cochrane.org/review-authors> (accessed 30 Jun 2015).
- Ali M, Schifano F, Robinson P, *et al*. Impact of community pharmacy diabetes monitoring and education programme on diabetes management: a randomized controlled study. *Diabet Med* 2012;29:e326–33.
- Fomos JA, Andres NF, Andres JC, *et al*. A pharmacotherapy follow-up program in patients with type-2 diabetes in community pharmacies in Spain. *Pharm World Sci* 2006;28:65–72.
- Paulos CP, Akesson Nygren CE, Celedon C, *et al*. Impact of a pharmaceutical care program in a community pharmacy on patients with dyslipidemia. *Ann Pharmacother* 2005;39:939–43.
- Correr CJ, Melchior AC, Fernandez-Llamos F, *et al*. Effects of a pharmacotherapy follow-up in community pharmacies on type 2 diabetes patients in Brazil. *Int J Clin Pharm* 2011;33:273–80.
- Zaragoza-Fernandez MP, Gastelurrutia MA, Cardero M, *et al*. Intensive two-month intervention on diet and lifestyle in uncontrolled hypertensive patients in a community pharmacy. *Latin Am J Pharm* 2012;31:727–33.
- Dhital R, Norman I, Whittlesea C, *et al*. The effectiveness of brief alcohol interventions delivered by community pharmacists: randomized controlled trial. *Addiction* 2015;110:1586–94.
- Watson MC, Stewart D. Screening and brief interventions for alcohol misuse delivered in the community pharmacy setting: a pilot study. Report for the Chief Scientist Office, 2011.
- Bauld L, Boyd KA, Briggs AH, *et al*. One-year outcomes and a cost-effectiveness analysis for smokers accessing group-based and pharmacy-led cessation services. *Nicotine Tob Res* 2011;13:135–45.
- Bock BC, Hudmon KS, Christian J, *et al*. A tailored intervention to support pharmacy-based counseling for smoking cessation. *Nicotine Tob Res* 2010;12:217–25.
- Burford O, Jiwa M, Carter O, *et al*. Internet-based photoaging within Australian pharmacies to promote smoking cessation: randomized controlled trial. *J Med Internet Res* 2013;15:e64.
- Costello MJ, Sproule B, Victor JC, *et al*. Effectiveness of pharmacist counseling combined with nicotine replacement therapy: a pragmatic randomized trial with 6,987 smokers. *Cancer Causes Control* 2011;22:167–80.
- Crealey GE, McElroy JC, Maguire TA, *et al*. Costs and effects associated with a community pharmacy-based smoking-cessation programme. *Pharmacoeconomics* 1998;14:323–33.
- Hoving C, Mudde AN, Dijk F, *et al*. Effectiveness of a smoking cessation intervention in Dutch pharmacies and general practices. *Health Educ* 2010;110:17–29.
- Howard-Pitney B, Killen JD, Fortmann SP. Quitting chew: results from a randomized trial using nicotine patches. *Exp Clin Psychopharmacol* 1999;7:362–71.
- Maguire TA, McElroy JC, Drummond A. A randomized controlled trial of a smoking cessation intervention based in community pharmacies. *Addiction* 2001;96:325–31.
- Mochizuki M, Hatsugaya M, Rokujoh E, *et al*. Randomized controlled study on the effectiveness of community pharmacists' advice for smoking cessation by Nicorette—evaluation at three months after initiation. *Yakugaku Zasshi* 2004;124:989–95.
- Sinclair HK, Bond CM, Lennox AS, *et al*. Training pharmacists and pharmacy assistants in the stage-of-change model of smoking cessation: a randomised controlled trial in Scotland. *Tob Control* 1998;7:253–61.
- Vial RJ, Jones TE, Ruffin RE, *et al*. Smoking cessation program using nicotine patches linking hospital to the community. *J Pharm Pract Res* 2002;32:57–62.
- Sonderskov J, Olsen J, Sabroe S, *et al*. Nicotine patches in smoking cessation: a randomized trial among over-the-counter customers in Denmark. *Am J Epidemiol* 1997;145:309–18.



41. Ahrens RA, Hower M, Best AM. Effects of weight reduction interventions by community pharmacists. *J Am Pharm Assoc* 2003;43:583–9.
42. Bush J, Langle CA, Patel A, *et al*. Evaluation of the Heart of Birmingham teaching Primary Care Trust (HoBtPCT) My Choice Weight Management Programme. Final Report; 2011.
43. Jolly K, Lewis A, Beach J, *et al*. Comparison of range of commercial or primary care led weight reduction programmes with minimal intervention control for weight loss in obesity: lighten Up randomised controlled trial. *BMJ* 2011;343:d6500.
44. Malone M, Alger-Mayer SA. Pharmacist intervention enhances adherence to orlistat therapy. *Ann Pharmacother* 2003;37:1598–602.
45. Phimarn W, Pianchana P, Limpikanchakovit P, *et al*. Thai community pharmacist involvement in weight management in primary care to improve patient's outcomes. *Int J Clin Pharm* 2013;35:1208–17.
46. Steed L, Kassavou A, Madurasinghe VW, *et al*. Community pharmacy interventions for health promotion: effects on professional practice and health outcomes. *Cochrane Database Syst Rev* 2014; (7):CD011207.

BMJ Open

Community pharmacy-delivered interventions for public health priorities: a systematic review of interventions for alcohol reduction, smoking cessation and weight management, including meta-analysis for smoking cessation

Tamara J Brown, Adam Todd, Claire O'Malley, Helen J Moore, Andrew K Husband, Clare Bamba, Adetayo Kasim, Falko F Sniehotta, Liz Steed, Sarah Smith, Lucie Nield and Carolyn D Summerbell

BMJ Open 2016 6:

doi: 10.1136/bmjopen-2015-009828

Updated information and services can be found at:
<http://bmjopen.bmj.com/content/6/2/e009828>

These include:

References

This article cites 31 articles, 11 of which you can access for free at:
<http://bmjopen.bmj.com/content/6/2/e009828#BIBL>

Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See:
<http://creativecommons.org/licenses/by/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Addiction](#) (105)
[Cardiovascular medicine](#) (536)
[Evidence based practice](#) (455)
[Public health](#) (1422)
[Smoking and tobacco](#) (180)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>